

Specific Procedure Post Treatment Instructions

Sinus Grafting Procedures

Treatment to the area of the sinus can cause slight discomfort, swelling and occasional bruising. It is advised to avoid bending over, nose blowing and suppressing a sneeze for 2 weeks.

- Slight nose bleeds are normal for up to 2 weeks.
- Small bone particles may be found in the mouth in the morning, they are normal.

Soft Tissue Gum Graft/ Connective Tissue Graft

During the normal course of healing, the tissue graft may change appearance and color. The color may appear white/gray/red during the healing period. Do NOT be alarmed by the appearance of the graft.

All-on-X/1Day Smile

Full mouth reconstructive surgery varies in the duration and extent of discomfort, swelling and occasional bruising. If treatment is completed on the upper jaw, avoid bending over, nose blowing and suppressing a sneeze for 2 weeks.

- Soft mushy diet to be followed for 6 weeks following procedure.
- Oral home care will not be implemented until instructed to do so.

LANAP/ LAPIP

It is extremely important not to dislodge blood clots that form in and around the gums. One of the most important results of laser surgery is the blood clots that form following surgery.

- Carefully avoid chewing food in the areas of the mouth where the laser has been used for at least one week following surgery.
- Do not be alarmed with any color changes or appearance of tissues following laser therapy. Tissues can be gray, yellow, red, blue, purple, “stringy” and reflect normal response to laser treatments.

Do NOT Rinse the first 24 Hours.

- It is OK to spit, rinse, and wash your mouth after 24 hours.

Do NOT brush or floss your teeth until instructed to do so.

- Try to keep your mouth as clean as possible to help the healing process. Brush, floss, and follow other home care measures in all the areas of your mouth except for the surgery area.
- Do not apply excessive tongue or cheek pressure to the surgery area.

Diet Instructions

- **The first three days**, follow **only** a liquid-like diet to allow healing. Anything that could be put into a blender to drink is ideal. The purpose of this is to protect the clot that is acting as a “band-aid” between the gums and the teeth. Do **not** drink through a straw, as this creates a vacuum in your mouth that can disturb the “band-aid”. Take daily vitamins.
- **Next four days after treatment**, foods with a “mushy” consistency such as

eggs, yogurt, cottage cheese, malts, ice cream, etc. Stay away from spicy and “chippy” foods. It is also important to maintain good fluid intake during this period.

- **Starting seven to ten days after treatment**, soft foods may be allowable. Soft foods have the consistency of pasta, fish, chicken or steamed vegetables. You may then, gradually add back your regular diet choices.

Please remember that even after ten days, healing is not complete. The first month following treatment you should continue to make smart food choices. Softer foods are better

Pinhole

Do Not brush, floss, touch at the surgical site. Rinse only with your lips apart, no vigorous swishing.

- Water-pik can be used only from the tongue-side (inside) surfaces.
- Tissue will be healing changes in color and appearance will occur over the next 6 weeks.
- Do not be alarmed about soft swelling or bruising the first week.
- Expect cold sensitivity on teeth for 6 weeks or longer.

Ice are at 10-minute intervals for the first 48 hours to minimize swelling.

- Pain can be controlled with one acetaminophen (Tylenol) and NSAID (Motrin or Advil) at the same time, subject to other instruction from the Doctor.
- Wear prescribed appliance 24 hours a day, if instructed. Do not clench or grind your teeth. Check with the Doctor about any other appliances (e.g., retainers, Invisalign, perio-protect, etc.)

Do not eat crunchy or sticky food or drink out of a straw.

Dressing Instructions

It is common for dressings to come fall off. If so, use a Q-tip dipped in Chlorhexidine Gluconate (Peridex) and gently swab the area. If the dressing becomes loose, simply remove it, do not try to push back in place, this will irritate the tissue.

AVOID FOR 3-5 DAYS POST-SURGERY

- Aspirin and other blood thinners. Please discuss this with your care provider.
- Vitamin E.
- Fish Oil.